

# Dosage Charts

## Acetaminophen (Tylenol)

		Old Infant's Concentrated Drops 80mg/0.8mL	New Infant's Concentrated Drops 160mg/5mL	Children's Suspension Liquid 160mg/5mL	Children's Soft Chewable Tablets 80mg each	Junior Strength Chewable Tablets 160mg each
Weight	Age	Dropperful (Use dropper)	Dropperful (Use dropper)	Teaspoon	Tablet	Tablet
	2-3 mos	1/2=(0.4 mL)	1/4=(1.25 mL)			
12-17 lbs	4-11 mos	1=(0.8 mL)	1/2=(2.5 mL)	1/2 tsp.		
18-23 lbs	12-23 mos	1 1/2=(0.8 + 0.4 mL)	3/4=(3.75 mL)	3/4 tsp.		
24-35 lbs	2-3 yrs	2=(0.8 + 0.8 mL)	1=(5 mL)	1 tsp.	2	
36-47 lbs	4-5 yrs			1 1/2 tsp.	3	
48-59 lbs	6-8 yrs			2 tsp.	4	2
60-71 lbs	9-10 yrs			2 1/2 tsp.	5	2 1/2
72-95 lbs	11 yrs			3 tsp.	6	3
96 lbs & over	12 yrs					4

**Please note:** One dose lasts 4 hours. No more than 5 doses in a 24 hour period.

If the child's weight and age do not correspond to the same row on the chart, choose the dosage corresponding to your child's weight.

## Ibuprofen (Motrin, Advil) Dosage

		Infant's Ibuprofen Concentrated Drops 50mg/1.25mL	Children's Ibuprofen Suspension 100mg/5mL	Children's Soft Chewable Tablets 50mg each	Junior Strength Chewable Tablets 100mg each	Junior Strength Ibuprofen Caplets 100mg
Weight	Age	Dropperful (Use dropper)	Teaspoon	Tablet	Tablet	Caplet
12-17 lbs	6-11 mos	1=(1.25 mL)				
18-23 lbs	12-23 mos	1 ½=(1.875 mL)				
24-35 lbs	2-3 yrs	2=(2.5 mL)	1 tsp.	2		
36-47 lbs	4-5 yrs		1½ tsp.	3		
48-59 lbs	6-8 yrs		2 tsp.	4	2	2
60-71 lbs	9-10 yrs		2 ½ tsp.	5	2 ½	2 ½
72-95 lbs	11 yrs		3 tsp.	6	3	3

**Please Note:** One dose lasts 6–8 hours. Not to be used in children under 6 months of age. No more than 5 doses in a 24 hour period. If the child's weight and age do not correspond to the same row on the chart, choose the dosage corresponding to your child's weight.

- Milliliter is abbreviated as mL; 5mL equals one teaspoon (tsp).
- Don't use household teaspoons, which can vary in size.
- Aspirin should not be used in children to treat fever or pain.