

Meningococcal B Vaccine:

What You Need to Know

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1. Why get vaccinated?

Meningococcal B vaccine can help protect against **meningococcal disease** caused by serogroup B. A different meningococcal vaccine is available that can help protect against serogroups A, C, W, and Y.

Meningococcal disease can cause meningitis (infection of the lining of the brain and spinal cord) and infections of the blood. Even when it is treated, meningococcal disease kills 10 to 15 infected people out of 100. And of those who survive, about 10 to 20 out of every 100 will suffer disabilities such as hearing loss, brain damage, kidney damage, loss of limbs, nervous system problems, or severe scars from skin grafts.

Meningococcal disease is rare and has declined in the United States since the 1990s. However, it is a severe disease with a significant risk of death or lasting disabilities in people who get it.

Anyone can get meningococcal disease. Certain people are at increased risk, including:

- Infants younger than one year old
- Adolescents and young adults 16 through 23 years old
- People with certain medical conditions that affect the immune system
- Microbiologists who routinely work with isolates of *N. meningitidis*, the bacteria that cause meningococcal disease
- People at risk because of an outbreak in their community

2. Meningococcal B vaccine

For best protection, more than 1 dose of a meningococcal B vaccine is needed. There are two meningococcal B vaccines available. The same vaccine must be used for all doses.

Meningococcal B vaccines are recommended for people 10 years or older who are at increased risk for serogroup B meningococcal disease, including:

- People at risk because of a serogroup B meningococcal disease outbreak
- Anyone whose spleen is damaged or has been removed, including people with sickle cell disease
- Anyone with a rare immune system condition called “complement component deficiency”
- Anyone taking a type of drug called a “complement inhibitor,” such as eculizumab (also called “Soliris”®) or ravulizumab (also called “Ultomiris”®)
- Microbiologists who routinely work with isolates of *N. meningitidis*

These vaccines may also be given to anyone 16 through 23 years old to provide short-term protection against most strains of serogroup B meningococcal disease, based on discussions between the patient and health care provider. The preferred age for vaccination is 16 through 18 years.



3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of meningococcal B vaccine**, or has any **severe, life-threatening allergies**
- Is **pregnant or breastfeeding**

In some cases, your health care provider may decide to postpone meningococcal B vaccination until a future visit.

Meningococcal B vaccination should be postponed for pregnant people unless the person is at increased risk and, after consultation with their health care provider, the benefits of vaccination are considered to outweigh the potential risks.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting meningococcal B vaccine.

Your health care provider can give you more information.

4. Risks of a vaccine reaction

- Soreness, redness, or swelling where the shot is given, tiredness, headache, muscle or joint pain, fever, or nausea can happen after meningococcal B vaccination. Some of these reactions occur in more than half of the people who receive the vaccine.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim.

7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at www.fda.gov/vaccines-blood-biologics/vaccines.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines.





BRIGHT FUTURES HANDOUT ► PARENT

15 THROUGH 17 YEAR VISITS

Here are some suggestions from Bright Futures experts that may be of value to your family.

✓ HOW YOUR FAMILY IS DOING

- Set aside time to be with your teen and really listen to her hopes and concerns.
- Support your teen in finding activities that interest him. Encourage your teen to help others in the community.
- Help your teen find and be a part of positive after-school activities and sports.
- Support your teen as she figures out ways to deal with stress, solve problems, and make decisions.
- Help your teen deal with conflict.
- If you are worried about your living or food situation, talk with us. Community agencies and programs such as SNAP can also provide information and assistance.

✓ YOUR GROWING AND CHANGING TEEN

- Make sure your teen visits the dentist at least twice a year.
- Give your teen a fluoride supplement if the dentist recommends it.
- Support your teen's healthy body weight and help him be a healthy eater.
 - Provide healthy foods.
 - Eat together as a family.
 - Be a role model.
- Help your teen get enough calcium with low-fat or fat-free milk, low-fat yogurt, and cheese.
- Encourage at least 1 hour of physical activity a day.
- Praise your teen when she does something well, not just when she looks good.

✓ YOUR TEEN'S FEELINGS

- If you are concerned that your teen is sad, depressed, nervous, irritable, hopeless, or angry, let us know.
- If you have questions about your teen's sexual development, you can always talk with us.

✓ HEALTHY BEHAVIOR CHOICES

- Know your teen's friends and their parents. Be aware of where your teen is and what he is doing at all times.
- Talk with your teen about your values and your expectations on drinking, drug use, tobacco use, driving, and sex.
- Praise your teen for healthy decisions about sex, tobacco, alcohol, and other drugs.
- Be a role model.
- Know your teen's friends and their activities together.
- Lock your liquor in a cabinet.
- Store prescription medications in a locked cabinet.
- Be there for your teen when she needs support or help in making healthy decisions about her behavior.

15 THROUGH 17 YEAR VISITS—PARENT



SAFETY

- Encourage safe and responsible driving habits.
 - Lap and shoulder seat belts should be used by everyone.
 - Limit the number of friends in the car and ask your teen to avoid driving at night.
 - Discuss with your teen how to avoid risky situations, who to call if your teen feels unsafe, and what you expect of your teen as a driver.
 - Do not tolerate drinking and driving.
- If it is necessary to keep a gun in your home, store it unloaded and locked with the ammunition locked separately from the gun.

Consistent with *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*, 4th Edition

For more information, go to <https://brightfutures.aap.org>.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances. Original handout included as part of the *Bright Futures Tool and Resource Kit*, 2nd Edition.

Inclusion in this handout does not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of the resources mentioned in this handout. Web site addresses are as current as possible but may change at any time.

The American Academy of Pediatrics (AAP) does not review or endorse any modifications made to this handout and in no event shall the AAP be liable for any such changes.

© 2019 American Academy of Pediatrics. All rights reserved.