



BRIGHT FUTURES HANDOUT ► PARENT

2½ YEAR VISIT

Here are some suggestions from Bright Futures experts that may be of value to your family.

✓ FAMILY ROUTINES

- Enjoy meals together as a family and always include your child.
- Have quiet evening and bedtime routines.
- Visit zoos, museums, and other places that help your child learn.
- Be active together as a family.
- Stay in touch with your friends. Do things outside your family.
- Make sure you agree within your family on how to support your child's growing independence, while maintaining consistent limits.

✓ LEARNING TO TALK AND COMMUNICATE

- Read books together every day. Reading aloud will help your child get ready for preschool.
- Take your child to the library and story times.
- Listen to your child carefully and repeat what she says using correct grammar.
- Give your child extra time to answer questions.
- Be patient. Your child may ask to read the same book again and again.

✓ GETTING ALONG WITH OTHERS

- Give your child chances to play with other toddlers. Supervise closely because your child may not be ready to share or play cooperatively.
- Offer your child and his friend multiple items that they may like. Children need choices to avoid battles.
- Give your child choices between 2 items your child prefers. More than 2 is too much for your child.
- Limit TV, tablet, or smartphone use to no more than 1 hour of high-quality programs each day. Be aware of what your child is watching.
- Consider making a family media plan. It helps you make rules for media use and balance screen time with other activities, including exercise.

✓ GETTING READY FOR PRESCHOOL

- Think about preschool or group child care for your child. If you need help selecting a program, we can give you information and resources.
- Visit a teachers' store or bookstore to look for books about preparing your child for school.
- Join a playgroup or make playdates.
- Make toilet training easier.
 - Dress your child in clothing that can easily be removed.
 - Place your child on the toilet every 1 to 2 hours.
 - Praise your child when he is successful.
- Try to develop a potty routine.
- Create a relaxed environment by reading or singing on the potty.

Helpful Resources: Family Media Use Plan: www.healthychildren.org/MediaUsePlan

Information About Car Safety Seats: www.safercar.gov/parents | Toll-free Auto Safety Hotline: 888-327-4236

2½ YEAR VISIT—PARENT

✓ SAFETY

- Make sure the car safety seat is installed correctly in the back seat. Keep the seat rear facing until your child reaches the highest weight or height allowed by the manufacturer. The harness straps should be snug against your child's chest.
- Everyone should wear a lap and shoulder seat belt in the car. Don't start the vehicle until everyone is buckled up.
- Never leave your child alone inside or outside your home, especially near cars or machinery.
- Have your child wear a helmet that fits properly when riding bikes and trikes or in a seat on adult bikes.
- Keep your child within arm's reach when she is near or in water.
- Empty buckets, play pools, and tubs when you are finished using them.
- When you go out, put a hat on your child, have her wear sun protection clothing, and apply sunscreen with SPF of 15 or higher on her exposed skin. Limit time outside when the sun is strongest (11:00 am–3:00 pm).
- Have working smoke and carbon monoxide alarms on every floor. Test them every month and change the batteries every year. Make a family escape plan in case of fire in your home.

WHAT TO EXPECT AT YOUR CHILD'S 3 YEAR VISIT

We will talk about

- Caring for your child, your family, and yourself
- Playing with other children
- Encouraging reading and talking
- Eating healthy and staying active as a family
- Keeping your child safe at home, outside, and in the car

Consistent with *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents, 4th Edition*

For more information, go to <https://brightfutures.aap.org>.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances. Original handout included as part of the *Bright Futures Tool and Resource Kit*, 2nd Edition.

Inclusion in this handout does not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of the resources mentioned in this handout. Web site addresses are as current as possible but may change at any time.

The American Academy of Pediatrics (AAP) does not review or endorse any modifications made to this handout and in no event shall the AAP be liable for any such changes.

© 2019 American Academy of Pediatrics. All rights reserved.



Dosage Charts

Milliliter is abbreviated as mL; 5mL equals one teaspoon (tsp)

Do not use household teaspoons, which can vary in size.

Aspirin should not be used in children to treat fever or pain.

483 East County Line Rd.
Hatboro, PA 19040

Keep all medications away from small children. In the event of an emergency please contact Poison Control: 1 (800) 222-1222

Acetaminophen (Tylenol)

2 MONTHS & OLDER ONLY

		Infant's Concentrated Drops 160 mg/ 5 mL	Children's Suspension Liquid 160 mg/ 5 mL	Children's Soft Chewable Tablets 80 mg each	Junior Strength Chewable Tablets 160 mg each
Weight	Age	Dropperful (Use dropper)	Teaspoon	Tablet	Tablet
6-11 lbs	2-3 Months	1.25 mL			
12-17 lbs	4-11 Months	2.5 mL	½ tsp (2.5mL)		
18-23 lbs	12-23 Months	3.75 mL	¾ tsp (3.75 mL)		
24-35 lbs	2-3 Years	5 mL	1 tsp (5 mL)	2	
36-47 lbs	4-5 Years		1 ½ tsp (7.5 mL)	3	
48-59 lbs	6-8 Years		2 tsp (10 mL)	4	2
60-71 lbs	9-10 Years		2 ½ tsp (12.5 mL)	5	2 ½
72-95 lbs	11 Years		3 tsp (15 mL)	6	3
96 lbs +	12 Years				4

Please note: One dose lasts 4 hours. No more than 5 doses in a 24 hour period. If the child's weight and age do not correspond to the same row on the chart, choose the dosage corresponding to your child's weight.

Ibuprofen (Motrin, Advil)

6 MONTHS & OLDER ONLY

		Infant's Concentrated Drops 50 mg/1.25 mL	Children's Suspension 100 mg/5 mL	Children's Soft Chewable Tablets 50 mg each	Junior Strength Chewable Tablets 100 mg each	Junior Strength Caplets 100 mg
Weight	Age	Dropperful (Use dropper)	Teaspoon	Tablet	Tablet	Caplet
12-17 lbs	6-11 Months	1.25 mL				
18-23 lbs	12-23 Months	1.875 mL				
24-35 lbs	2-3 Years	2.5 mL	1 tsp (5 mL)	2		
36-47 lbs	4-5 Years		1 ½ tsp (7.5 mL)	3		
48-59 lbs	9-10 Years		2 tsp (10 mL)	4	2	2
60-71 lbs	9-10 Years		2 ½ tsp (12.5 mL)	5	2 ½	2 ½
72-95 lbs	11 Years		3 tsp (15 mL)	6	3	3

Please Note: One dose lasts 6-8 hours. Not to be used in children under 6 months of age. No more than 5 doses in a 24 hour period. If the child's weight and age do not correspond to the same row on the chart, choose the dosage corresponding to your child's weight.