

Your Child's First Vaccines: What You Need to Know

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

The vaccines included on this statement are likely to be given at the same time during infancy and early childhood. There are separate Vaccine Information Statements for other vaccines that are also routinely recommended for young children (measles, mumps, rubella, varicella, rotavirus, influenza, and hepatitis A).

Your child is getting these vaccines today:

DTaP Hib Hepatitis B Polio PCV13

(Provider: Check appropriate boxes.)

1. Why get vaccinated?

Vaccines can prevent disease. Childhood vaccination is essential because it helps provide immunity before children are exposed to potentially life-threatening diseases.

Diphtheria, tetanus, and pertussis (DTaP)

- **Diphtheria (D)** can lead to difficulty breathing, heart failure, paralysis, or death.
- **Tetanus (T)** causes painful stiffening of the muscles. Tetanus can lead to serious health problems, including being unable to open the mouth, having trouble swallowing and breathing, or death.
- **Pertussis (aP)**, also known as “whooping cough,” can cause uncontrollable, violent coughing that makes it hard to breathe, eat, or drink. Pertussis can be extremely serious especially in babies and young children, causing pneumonia, convulsions, brain damage, or death. In teens and adults, it can cause weight loss, loss of bladder control, passing out, and rib fractures from severe coughing.

Hib (*Haemophilus influenzae* type b) disease

Haemophilus influenzae type b can cause many different kinds of infections. These infections usually affect children under 5 years of age but can also affect adults with certain medical conditions. Hib bacteria can cause mild illness, such as ear infections

or bronchitis, or they can cause severe illness, such as infections of the blood. Severe Hib infection, also called “invasive Hib disease,” requires treatment in a hospital and can sometimes result in death.

Hepatitis B

Hepatitis B is a liver disease that can cause mild illness lasting a few weeks, or it can lead to a serious, lifelong illness. Acute hepatitis B infection is a short-term illness that can lead to fever, fatigue, loss of appetite, nausea, vomiting, jaundice (yellow skin or eyes, dark urine, clay-colored bowel movements), and pain in the muscles, joints, and stomach. Chronic hepatitis B infection is a long-term illness that occurs when the hepatitis B virus remains in a person's body. Most people who go on to develop chronic hepatitis B do not have symptoms, but it is still very serious and can lead to liver damage (cirrhosis), liver cancer, and death.

Polio

Polio (or poliomyelitis) is a disabling and life-threatening disease caused by poliovirus, which can infect a person's spinal cord, leading to paralysis. Most people infected with poliovirus have no symptoms, and many recover without complications. Some people will experience sore throat, fever, tiredness, nausea, headache, or stomach pain.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

A smaller group of people will develop more serious symptoms: paresthesia (feeling of pins and needles in the legs), meningitis (infection of the covering of the spinal cord and/or brain), or paralysis (can't move parts of the body) or weakness in the arms, legs, or both. Paralysis can lead to permanent disability and death.

Pneumococcal disease

Pneumococcal disease refers to any illness caused by pneumococcal bacteria. These bacteria can cause many types of illnesses, including pneumonia, which is an infection of the lungs. Besides pneumonia, pneumococcal bacteria can also cause ear infections, sinus infections, meningitis (infection of the tissue covering the brain and spinal cord), and bacteremia (infection of the blood). Most pneumococcal infections are mild. However, some can result in long-term problems, such as brain damage or hearing loss. Meningitis, bacteremia, and pneumonia caused by pneumococcal disease can be fatal.

2. DTaP, Hib, hepatitis B, polio, and pneumococcal conjugate vaccines

Infants and children usually need:

- 5 doses of **diphtheria, tetanus, and acellular pertussis vaccine (DTaP)**
- 3 or 4 doses of **Hib vaccine**
- 3 doses of **hepatitis B vaccine**
- 4 doses of **polio vaccine**
- 4 doses of **pneumococcal conjugate vaccine (PCV13)**

Some children might need fewer or more than the usual number of doses of some vaccines to be fully protected because of their age at vaccination or other circumstances.

Older children, adolescents, and adults with certain health conditions or other risk factors might also be recommended to receive 1 or more doses of some of these vaccines.

These vaccines may be given as stand-alone vaccines, or as part of a combination vaccine (a type of vaccine that combines more than one vaccine together into one shot).

3. Talk with your health care provider

Tell your vaccination provider if the child getting the vaccine:

For all of these vaccines:

- Has had an **allergic reaction after a previous dose of the vaccine**, or has any **severe, life-threatening allergies**

For DTaP:

- Has had an **allergic reaction after a previous dose of any vaccine that protects against tetanus, diphtheria, or pertussis**
- Has had a **coma, decreased level of consciousness, or prolonged seizures within 7 days after a previous dose of any pertussis vaccine (DTP or DTaP)**
- Has **seizures or another nervous system problem**
- Has ever had **Guillain-Barré Syndrome** (also called “GBS”)
- Has had **severe pain or swelling after a previous dose of any vaccine that protects against tetanus or diphtheria**

For PCV13:

- Has had an **allergic reaction after a previous dose of PCV13, to an earlier pneumococcal conjugate vaccine known as PCV7, or to any vaccine containing diphtheria toxoid** (for example, DTaP)

In some cases, your child's health care provider may decide to postpone vaccination until a future visit.

Children with minor illnesses, such as a cold, may be vaccinated. Children who are moderately or severely ill should usually wait until they recover before being vaccinated.

Your child's health care provider can give you more information.

4. Risks of a vaccine reaction

For all of these vaccines:

- Soreness, redness, swelling, warmth, pain, or tenderness where the shot is given can happen after vaccination.

For DTaP vaccine, Hib vaccine, hepatitis B vaccine, and PCV13:

- Fever can happen after vaccination.

For DTaP vaccine:

- Fussiness, feeling tired, loss of appetite, and vomiting sometimes happen after DTaP vaccination.
- More serious reactions, such as seizures, non-stop crying for 3 hours or more, or high fever (over 105°F) after DTaP vaccination happen much less often. Rarely, vaccination is followed by swelling of the entire arm or leg, especially in older children when they receive their fourth or fifth dose.

For PCV13:

- Loss of appetite, fussiness (irritability), feeling tired, headache, and chills can happen after PCV13 vaccination.
- Young children may be at increased risk for seizures caused by fever after PCV13 if it is administered at the same time as inactivated influenza vaccine. Ask your health care provider for more information.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim.

7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at www.fda.gov/vaccines-blood-biologics/vaccines.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines.



Rotavirus Vaccine:

What You Need to Know

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1. Why get vaccinated?

Rotavirus vaccine can prevent rotavirus disease.

Rotavirus commonly causes severe, watery diarrhea, mostly in babies and young children. Vomiting and fever are also common in babies with rotavirus. Children may become dehydrated and need to be hospitalized and can even die.

2. Rotavirus vaccine

Rotavirus vaccine is administered by putting drops in the child's mouth. Babies should get 2 or 3 doses of rotavirus vaccine, depending on the brand of vaccine used.

- The first dose must be administered before 15 weeks of age.
- The last dose must be administered by 8 months of age.

Almost all babies who get rotavirus vaccine will be protected from severe rotavirus diarrhea.

Another virus called "porcine circovirus" can be found in one brand of rotavirus vaccine (Rotarix). This virus does not infect people, and there is no known safety risk.

Rotavirus vaccine may be given at the same time as other vaccines.

3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of rotavirus vaccine**, or has any **severe, life-threatening allergies**
- Has a **weakened immune system**
- Has **severe combined immunodeficiency (SCID)**
- Has had a type of bowel blockage called "**intussusception**"

In some cases, your child's health care provider may decide to postpone rotavirus vaccination until a future visit.

Infants with minor illnesses, such as a cold, may be vaccinated. Infants who are moderately or severely ill should usually wait until they recover before getting rotavirus vaccine.

Your child's health care provider can give you more information.

4. Risks of a vaccine reaction

- Irritability or mild, temporary diarrhea or vomiting can happen after rotavirus vaccine.

Intussusception is a type of bowel blockage that is treated in a hospital and could require surgery. It happens naturally in some infants every year in the United States, and usually there is no known reason for it. There is also a small risk of intussusception from rotavirus vaccination, usually within a week after the first or second vaccine dose. This additional risk is estimated to range from about 1 in 20,000 U.S. infants to 1 in 100,000 U.S. infants who get rotavirus vaccine. Your health care provider can give you more information.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

5. What if there is a serious problem?

For intussusception, look for signs of stomach pain along with severe crying. Early on, these episodes could last just a few minutes and come and go several times in an hour. Babies might pull their legs up to their chest. Your baby might also vomit several times or have blood in the stool, or could appear weak or very irritable. These signs would usually happen during the first week after the first or second dose of rotavirus vaccine, but look for them any time after vaccination. If you think your baby has intussusception, contact a health care provider right away. If you can't reach your health care provider, take your baby to a hospital. Tell them when your baby got rotavirus vaccine.

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim.

7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at www.fda.gov/vaccines-blood-biologics/vaccines.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636** (**1-800-CDC-INFO**) or
 - Visit CDC's website at www.cdc.gov/vaccines.





BRIGHT FUTURES HANDOUT ► PARENT

2 MONTH VISIT

Here are some suggestions from Bright Futures experts that may be of value to your family.

✓ HOW YOUR FAMILY IS DOING

- If you are worried about your living or food situation, talk with us. Community agencies and programs such as WIC and SNAP can also provide information and assistance.
- Find ways to spend time with your partner. Keep in touch with family and friends.
- Find safe, loving child care for your baby. You can ask us for help.
- Know that it is normal to feel sad about leaving your baby with a caregiver or putting him into child care.

✓ HOW YOU ARE FEELING

- Take care of yourself so you have the energy to care for your baby.
- Talk with me or call for help if you feel sad or very tired for more than a few days.
- Find small but safe ways for your other children to help with the baby, such as bringing you things you need or holding the baby's hand.
- Spend special time with each child reading, talking, and doing things together.

✓ FEEDING YOUR BABY

- Feed your baby only breast milk or iron-fortified formula until she is about 6 months old.
- Avoid feeding your baby solid foods, juice, and water until she is about 6 months old.
- Feed your baby when you see signs of hunger. Look for her to
 - Put her hand to her mouth.
 - Suck, root, and fuss.
- Stop feeding when you see signs your baby is full. You can tell when she
 - Turns away
 - Closes her mouth
 - Relaxes her arms and hands
- Burp your baby during natural feeding breaks.

If Breastfeeding

- Feed your baby on demand. Expect to breastfeed 8 to 12 times in 24 hours.
- Give your baby vitamin D drops (400 IU a day).
- Continue to take your prenatal vitamin with iron.
- Eat a healthy diet.
- Plan for pumping and storing breast milk. Let us know if you need help.
 - If you pump, be sure to store your milk properly so it stays safe for your baby. If you have questions, ask us.

If Formula Feeding

- Feed your baby on demand. Expect her to eat about 6 to 8 times each day, or 26 to 28 oz of formula per day.
- Make sure to prepare, heat, and store the formula safely. If you need help, ask us.
- Hold your baby so you can look at each other when you feed her.
- Always hold the bottle. Never prop it.

✓ YOUR GROWING BABY

- Have simple routines each day for bathing, feeding, sleeping, and playing.
- Hold, talk to, cuddle, read to, sing to, and play often with your baby. This helps you connect with and relate to your baby.
- Learn what your baby does and does not like.
- Develop a schedule for naps and bedtime. Put him to bed awake but drowsy so he learns to fall asleep on his own.
- Don't have a TV on in the background or use a TV or other digital media to calm your baby.
- Put your baby on his tummy for short periods of playtime. Don't leave him alone during tummy time or allow him to sleep on his tummy.
- Notice what helps calm your baby, such as a pacifier, his fingers, or his thumb. Stroking, talking, rocking, or going for walks may also work.
- *Never hit or shake your baby.*

Helpful Resources:

Information About Car Safety Seats: www.safercar.gov/parents | Toll-free Auto Safety Hotline: 888-327-4236

2 MONTH VISIT—PARENT



SAFETY

- Use a rear-facing-only car safety seat in the back seat of all vehicles.
- Never put your baby in the front seat of a vehicle that has a passenger airbag.
- Your baby's safety depends on you. Always wear your lap and shoulder seat belt. Never drive after drinking alcohol or using drugs. Never text or use a cell phone while driving.
- Always put your baby to sleep on her back in her own crib, not your bed.
 - Your baby should sleep in your room until she is at least 6 months old.
 - Make sure your baby's crib or sleep surface meets the most recent safety guidelines.
- If you choose to use a mesh playpen, get one made after February 28, 2013.
- Swaddling should not be used after 2 months of age.
- Prevent scalds or burns. Don't drink hot liquids while holding your baby.
- Prevent tap water burns. Set the water heater so the temperature at the faucet is at or below 120°F /49°C.
- Keep a hand on your baby when dressing or changing her on a changing table, couch, or bed.
- Never leave your baby alone in bathwater, even in a bath seat or ring.

WHAT TO EXPECT AT YOUR BABY'S 4 MONTH VISIT

We will talk about

- Caring for your baby, your family, and yourself
- Creating routines and spending time with your baby
- Keeping teeth healthy
- Feeding your baby
- Keeping your baby safe at home and in the car

Consistent with *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*, 4th Edition

For more information, go to <https://brightfutures.aap.org>.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances. Original handout included as part of the *Bright Futures Tool and Resource Kit*, 2nd Edition.

Inclusion in this handout does not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of the resources mentioned in this handout. Web site addresses are as current as possible but may change at any time.

The American Academy of Pediatrics (AAP) does not review or endorse any modifications made to this handout and in no event shall the AAP be liable for any such changes.

© 2019 American Academy of Pediatrics. All rights reserved.



Dosage Charts

Milliliter is abbreviated as mL; 5mL equals one teaspoon (tsp)

Do not use household teaspoons, which can vary in size.

Aspirin should not be used in children to treat fever or pain.

483 East County Line Rd.
Hatboro, PA 19040

Keep all medications away from small children. In the event of an emergency please contact Poison Control: 1 (800) 222-1222

Acetaminophen (Tylenol)

2 MONTHS & OLDER ONLY

		Infant's Concentrated Drops 160 mg/ 5 mL	Children's Suspension Liquid 160 mg/ 5 mL	Children's Soft Chewable Tablets 80 mg each	Junior Strength Chewable Tablets 160 mg each
Weight	Age	Dropperful (Use dropper)	Teaspoon	Tablet	Tablet
6-11 lbs	2-3 Months	1.25 mL			
12-17 lbs	4-11 Months	2.5 mL	½ tsp (2.5mL)		
18-23 lbs	12-23 Months	3.75 mL	¾ tsp (3.75 mL)		
24-35 lbs	2-3 Years	5 mL	1 tsp (5 mL)	2	
36-47 lbs	4-5 Years		1 ½ tsp (7.5 mL)	3	
48-59 lbs	6-8 Years		2 tsp (10 mL)	4	2
60-71 lbs	9-10 Years		2 ½ tsp (12.5 mL)	5	2 ½
72-95 lbs	11 Years		3 tsp (15 mL)	6	3
96 lbs +	12 Years				4

Please note: One dose lasts 4 hours. No more than 5 doses in a 24 hour period. If the child's weight and age do not correspond to the same row on the chart, choose the dosage corresponding to your child's weight.

Ibuprofen (Motrin, Advil)

6 MONTHS & OLDER ONLY

		Infant's Concentrated Drops 50 mg/1.25 mL	Children's Suspension 100 mg/5 mL	Children's Soft Chewable Tablets 50 mg each	Junior Strength Chewable Tablets 100 mg each	Junior Strength Caplets 100 mg
Weight	Age	Dropperful (Use dropper)	Teaspoon	Tablet	Tablet	Caplet
12-17 lbs	6-11 Months	1.25 mL				
18-23 lbs	12-23 Months	1.875 mL				
24-35 lbs	2-3 Years	2.5 mL	1 tsp (5 mL)	2		
36-47 lbs	4-5 Years		1 ½ tsp (7.5 mL)	3		
48-59 lbs	9-10 Years		2 tsp (10 mL)	4	2	2
60-71 lbs	9-10 Years		2 ½ tsp (12.5 mL)	5	2 ½	2 ½
72-95 lbs	11 Years		3 tsp (15 mL)	6	3	3

Please Note: One dose lasts 6-8 hours. Not to be used in children under 6 months of age. No more than 5 doses in a 24 hour period. If the child's weight and age do not correspond to the same row on the chart, choose the dosage corresponding to your child's weight.